Temperature Profiles and the Effects of Field Environments on Germination of Silver Sagebrush

Artemisia cana ssp. *cana*

Photo by James T Romo

JAMES T ROMO AND JAMES A YOUNG

ABSTRACT

In the laboratory, silver sagebrush (*Artemisia cana* Pursh ssp. *cana* [Asteraceae]) seeds germinated with increasing temperatures similar to those of spring and early summer or with cooling temperatures that occur in late summer or autumn. However, exposing seeds to similar temperatures under field environments reduced germination, with greater reductions occurring when seeds were put in the field in autumn (–36% to –81%) compared to spring (–18% to –24%). Germination of seeds placed in the field in autumn declined abruptly as daily temperatures rose in early spring, however, a gradual decline in germination was noted for seeds put in the field in spring. Nearly all non-germinating seeds were infected with fungi and were non-viable, indicating that recruitment of seedlings may be limited by loss of viability of seeds in the soil. Despite the fact that viability of seeds declined with exposure to field environments, 19% to 82% of the seeds remained viable, with greater viability in spring- than autumn-placed seeds. Seeding silver sagebrush in spring is recommended to maintain viability and high germination of seeds.

KEY WORDS: *Artemisia cana* ssp. *cana*, ecological restoration, seed deterioration, seed germination, seedbed ecology

NOMENCLATURE: Looman and Best (1979)

Shrubs are important in many aspects of the structure and functioning of ecosystems (Miller 1987; Allen 1988; Call and Roundy 1991; Pyke and Archer 1991). For these reasons, restoration specialists are interested in planting silver sagebrush (*Artemisia cana* Pursh. ssp. *cana* [Asteraceae]) on drastically disturbed sites in the Northern Mixed Prairie. Northern Mixed Prairie lays in the northern portion of the Great Plains from southeastern Wyoming and northwestern Nebraska through the Dakotas and Montana to southern Alberta, Saskatchewan, and southwestern Manitoba (Launenroth and Milchunas 1992). The Northern Great Plains encompasses the region between the foothills of the Rocky Mountains in

TABLE 1

Storage locations (seed treatment), the time of the year when silver sagebrush seeds were placed in the field, and the number of seed groups in 3 y of study

Years of study	Storage location (seed treatment)		
	Laboratory		Field
	Time of placement		
	Autumn	Autumn	Spring
	(Number of seed groups)		
1994—1995	60	60	60
1995—1996	60	60	60
1998—1999	80	80	—

the west, approximately the 100th meridian on the east, the North Platte River in Wyoming and Nebraska in the south, and southern Alberta, Saskatchewan, and Manitoba in the north (Barker and Whitman 1988). Northern Mixed Prairie is characterized by a number of mid- and short-grasses (Barker and Whitman 1988; Coupland 1992) of the genera *Agropyron*, *Stipa*, *Koeleria*, and *Bouteloua* (Coupland 1992).

With the exception of some outlying populations, distribution of silver sagebrush largely corresponds with the Northern Mixed Prairie and the Northern Great Plains (Harvey 1981). This long-lived, rhizomatous shrub occupies early to late successional communities on a variety of soils over about 14 million ha (35 million acres) (Beetle 1960). In Canada, silver sagebrush is most abundant in the *Bouteloua-Stipa* faciation, and is less common in *Stipa-Bouteloua*, *Stipa-Bouteloua-Agropyron*, *Bouteloua-Agropyron*, and *Stipa-Agropyron* faciatiions (Coupland 1950, 1961). Silver sagebrush can be scattered throughout the grassland matrix (Rowe and Coupland 1984) or it can be a dominant in sandhill prairies (Hulett and others 1966), coulees (Lawrence and Romo 1994), or on flats that are periodically flooded in valleys and along streams (Hanson and Whitman 1938).

Silver sagebrush establishes primarily by sprouting from rhizomes (Wambolt and others 1990) and establishment of plants from seeding is generally limited (Eddleman 1980; Walton 1984; Romo and Grilz 2001). Burial of seeds too deep in the soil (Walton 1984), predation on seeds (Romo and Grilz 2001), very specific safe site requirements for germination (Romo and Grilz 2001), soil water limitations (Walton 1984), competition from surrounding vegetation (Walton 1984; Romo and Grilz 2001), death of seedlings caused by freezing temperatures (Hou and Romo 1998b), poorly adapted ecotypes (Meyer and Monsen 1992), and adverse environmental conditions (Walton 1984) may limit establishment of silver sagebrush from seeds.

Schuman and others (1998) speculated that seeds of Wyoming big sagebrush (*Artemisia tridentata* ssp. *wyomingensis* Beetle & Young) may enter dormancy and may remain viable in the soil. Fringed sagebrush (*Artemisia frigida* Willd), a suffrutescent commonly associated with silver sagebrush, develops a persistent seedbank (Bai and Romo 1997) and shows no seasonal pattern of seed dormancy (Bai and Romo 1994). Exposing seeds to -23°C (-9°F) for 30 d has no affect on germination of silver sagebrush (Walton 1984). Silver sagebrush seeds are not known to exhibit dormancy (Walton 1984) and other sagebrush species germinate over a wide range of temperatures (Harniss and McDonough 1976; Young and others 1991; Bai and others 1995).

Reductions in germination for seeds with high water content have been noted for many species (Hegarty 1978; Roos 1986). At 20°C , reduced seed viability was observed at 60 d when water content was around 5% in Wyoming big sagebrush and 7% in mountain big sagebrush (*Artemisia tridentata* Nutt. ssp. *vaseyana* [Rydb.] Beetle) (Welch 1996). By comparison, germination of Wyoming sagebrush was unaffected by maintaining seeds for up to 15 d at water contents of 40% to 60% at temperatures of 2 to 15°C (Bai and others 1997). Viability of mountain big sagebrush seeds declines faster when stored in widely fluctuating temperatures than at a constant, low temperature (Welch and others 1996). Seeds of fringed sagebrush remain viable for at least 3.5 to 5 y in the field (Bai and Romo 1994). The effects on germination of exposing silver sagebrush seeds to varying temperature and moisture conditions have not, however, been determined. It is possible that silver sagebrush seeds enter dormancy, lose viability, or are unaffected when exposed to environmental conditions in the field.

Our study objectives were to determine: 1) effects of exposing seeds to environmental conditions in the field on germination and viability of seeds of silver sagebrush, and 2) germination of seeds under constant and alternating temperatures in the laboratory. Basic requirements and characteristics of germination must be determined to increase establishment of silver sagebrush during restoration. Such information will aid restoration specialists in selecting times for seeding silver sagebrush to produce the most consistent success in establishing this shrub on drastically disturbed sites.

MATERIALS AND METHODS

Seed Source

Silver sagebrush seeds (achenes) were collected in mid-October 1994, 1995, and 1998 near Outlook, Saskatchewan (Lat $51^{\circ} 29'\text{N}$, Long $107^{\circ} 03'\text{W}$,

elevation 518 m [567 ft]). The collection site, a naturally vegetated roadcut, was dominated by silver sagebrush, western wheatgrass (*Agropyron smithii* Rydb. [Poaceae]), and needle-and-thread (*Stipa comata* Trin. & Rupr. [Poaceae]). Inflorescences of silver sagebrush were cut from plants, placed in large paper bags, and air-dried in a laboratory at room temperature for about 5 d. Inflorescences were rubbed by hand to remove seeds, and seeds were cleaned with a Clipper desktop thresher (Seedburo Equipment Company, Chicago, Illinois). Cleaned seeds were stored in paper envelopes in darkness at 5 °C (41 °F) until use.

Exposure of Seeds to Field Conditions

In 1994, 1995, and 1998, groups of 50 seeds were counted and put in paper envelopes wherein seeds could be exposed to ambient temperature and moisture conditions in the field. Seeds were stored in darkness at 5 °C (41 °F) until placed in the field. Seed-containing envelopes were put in the field at the University of Saskatchewan, Kernen Research Farm 1 km (0.6 miles) east of Saskatoon on 18 November 1994, 2 April 1995, 7 November 1995, 10 April 1996, and 17 November 1998 (Table 1). In 1994–1995, autumn-placed seeds were in the field for a total of 199 d while spring-placed seeds were exposed to the field environment for 64 d. Seeds were in the field for 211 and 57 d for autumn- and spring-placed seeds, respectively, in 1995–1996. Seeds were in the field for 216 d in 1998–1999. Envelopes were placed in two 50 by 80-cm (19.7 by 31.5-in) nylon, mesh bags and bags were secured to the ground with spikes. Envelopes were spread flat on the soil surface within the bags.

From 2 April to 5 June 1995, 10 April through 6 June 1996, and 23 January 1999 to 21 June 1999, 8 paper envelopes containing seeds placed in the field in autumn or spring were taken from the field at about 2- to 4-wk intervals. Eight replicates of seeds stored at 5 °C (41 °F) in the laboratory were processed at the same times. Water content of seeds, expressed on a fresh weight basis, was determined for 4 replicates per treatment by weighing seeds after blotting dry with tissue paper, drying at 80 °C (176 °F) for 48 h, and re-weighing. For laboratory-stored seeds the weight per 50 seeds on a fresh weight basis averaged 36 mg ($s_x = \pm 1$) in 1994, 34 mg ($s_x = \pm 1$) in 1995, and 31 mg ($s_x = \pm 1$) in 1998. Another 4 replicates of each treatment were prepared for testing germination by placing seeds in Petri dishes containing 1 layer of #4 Whatman filter paper moistened with distilled water. Closed Petri dishes were enclosed in clear plastic bags and incubated for 20 d at 10 °C (50 °F) with 12 h darkness and 12 h light ($220 \mu\text{mol}/(\text{m}^2 \cdot \text{s})$). Previous trials in our laboratory indicated complete

germination of silver sagebrush seeds when incubated under these conditions. Germinated seeds were counted and removed at 4-d intervals; distilled water was added to the filter paper if needed. In 1998–1999, the number of seeds that did not germinate but were infected with fungi was recorded. A seed was considered infected if fungi could be observed on it with the naked eye. Since it was not known whether these seeds were dormant or non-viable, they were transferred to a new Petri dish and soaked in a 2% solution of tetrazolium chloride for 48 h (Grabe 1972). Seeds were dissected and deemed viable if the cotyledons and radicle were stained dark red-purple by the tetrazolium chloride.

Figure 1 • Daily maximum and minimum temperatures to which seeds of silver sagebrush were exposed when put in the field on 18 November 1994 or 2 April 1995, 7 November 1995 or 10 April 1996, and 17 November 1998.

A Campbell Scientific 21X data logger (Campbell Scientific Canada Corporation, Edmonton, AB) with temperature sensors was placed in the field at the same time seeds were put out in autumn. Two temperature sensors were placed in paper envelopes and daily minimum and maximum temperatures were recorded through the duration of the experiments. Precipitation was recorded continuously at the Kernen Research Farm weather station about 100 m (109 ft) from where seeds were placed.

TABLE 2

Water content of seeds and total germination for silver sagebrush during the seed retrieval test in 1994–1995

Seed treatment (location and time of placement)	Date of retrieval in 1994–1995						
	18 Nov	2 Apr	15 Apr	27 Apr	8 May	23 May	5 Jun
	Water content of seeds (% fresh weight)						
Laboratory	23	13	11	9	9	6	11
Field—Autumn	19	36	40	23	37	16	36
Field—Spring	12	9	40	19	28	11	23
Standard error (date of germination X time of placement interaction)							2.4
	Total germination (%)						
Laboratory	95	89	94	87	91	94	96
Field—Autumn	92	83	75	68	63	55	59
Field—Spring	98	95	92	98	91	88	80
Standard error (date of germination X time of placement interaction)							3.9

In 1994–1995, seeds placed in the field in autumn were exposed to temperatures ranging from -23 to 54 °C (-9 to 129 °F) while seeds put out in spring experienced temperatures between -20 and 54 °C (-4 to 129 °F) (Figure 1). Diurnal variation in temperatures ranged from 1 to 51 °C (34 to 124 °F) for seeds put in the field in autumn, and from 6 to 51 °C (43 to 124 °F) for seeds put in the field in spring. Seeds were covered by snow (77 mm [3.0 in] of precipitation) after being put in the field in November 1994 through March 1995. In 1995, 34 mm (1.3 in) of precipitation were received in April, and 15 mm (0.6 in) were received in May.

During 1995–1996, seeds placed in the field were exposed to temperatures between -28 and 43 °C (-18 to 109 °F) in autumn and -9 to 43 °C (16 to 109 °F) in spring (Figure 1). Diurnal variation ranged from about 1 to 43 °C (34 to 109 °F) in autumn and 6 to 43 °C (43 to 109 °F) in spring. Seeds were covered by snow from November 1995 through late March 1996, which provided 76 mm (3.0 in) precipitation. Monthly precipitation totaled 30 mm (1.2 in) in April, 59 mm (2.3 in) in May, and 101 mm (4.0 in) in June.

Seeds placed in the field in 1998–1999 were exposed to temperatures ranging from -19 to 51 °C (-2 to 124 °F), with diurnal variation from 0 to 38 °C (32 to 100 °F) (Figure 1). Seeds were covered with snow from December through late March, which contained 58 mm (2.3 in) precipitation. Precipitation totaled 15 mm (0.6 in) in April, 144 mm (5.7 in) in May, and 59 mm (2.3 in) in June.

In the 1994–1995 and 1995–1996 experiments, the experimental design was a randomized-complete-block with placement of seeds (seed treatment) in the field in autumn and spring and

laboratory storage, and dates of germination applied factorially (Snedecor and Cochran 1980). Main effects and their interactions on water content of seeds and total germination were tested with analysis of variance (Snedecor and Cochran 1980). Linear, quadratic, and cubic contrasts were performed on means of total germination (Petersen 1985). These contrasts compared mean germination over time for seeds stored in the laboratory with that of seeds put in the field in autumn and spring, and for seeds placed in the field in autumn versus placement in the spring. Data from each year were analyzed separately because of different and an unequal number of sample dates. In the 1998–1999 experiment, the seed treatments were factorially applied in combination with dates of germination in randomized-complete-block design. Main effects and their interacting effects on water content of seeds, total germination, percentage of seeds infected with fungi, and seeds staining with tetrazolium chloride were tested with analysis of variance. In all cases statistical significance was presumed at $P < 0.05$.

Germination of Seeds Under Constant and Alternate Temperatures in the Laboratory

Seeds were sent to the USDA Agriculture Research Service Laboratory in Reno, Nevada, in 1994, 1995, and 1998, where germination was investigated in 55 constant and alternating temperatures (Evans and others 1982) about 3 mo after collection. A randomized-complete-block-design with 25 seeds in each of 4 replicates was used. Seeds were incubated in dark germinators in closed Petri dishes on 1-mm thick germination paper that was kept moist with water. Constant temperature regimes were 0 , 2 , 5 , and 5 °C (32 , 36 , 41 and 9 °F) incre-

ments through 40 °C (104 °F). Alternating temperature regimes consisted of a 16-h cold period and an 8-h warm period, at all possible higher temperatures each 24-h interval. For example, a 2 °C (36 °F) cold period was alternated with 5, 10, 15, 20, 25, 30, 35, or 40 °C (41, 50, 59, 68, 77, 86, 95, 104 °F) warm period, whereas a 30 °C (86 °F) cold period was alternated with a warm period of 35 or 40 °C (95 or 104 °F). Germination of seeds was recorded after 4 wk of incubation. Four wk of incubation of wildland species is standard protocol in this laboratory, thus it was used in these germination tests. Response surfaces were developed for germination of each collection using multiple regression analysis (Evans and others 1982).

RESULTS

Exposure of Seeds to Laboratory and Field Environments

1994–1995 and 1995–1996

In both years the interacting effects of seed treatment and the date of germination affected water content of seeds and total germination ($P = 0.000$ to 0.002). Water content of seeds varied widely among dates, and it averaged 30% for autumn-placed seeds, 20% for spring-placed seeds, and 12% for seeds stored in the laboratory in 1994–1995 (Table 2). In 1995–1996, water content of seeds in the field averaged 23% and 18% when put out in autumn and spring, respectively, while that of laboratory-stored seeds averaged 10% (Table 3).

More seeds germinated when stored in the laboratory than if exposed to field environments, and

germination was greater for seeds placed in the field in spring compared to those put out in autumn (Tables 2 and 3). Contrasts of means indicated total germination changed through time in unique ways for laboratory-stored seeds versus seeds placed in the field (Table 4). The field environment affected total germination of autumn- and spring-placed seeds differently as indicated by changes in germination over time (Table 4). From the first to the last date, germination declined 36% for autumn-placed seeds in 1994–1995 and 41% in 1995–1996. Over the same period, an 18% reduction occurred for total germination in 1994–1995 and 24% in 1995–1996 for seeds put in the field in spring. In both years the decline in germination in April for seeds placed in the field in autumn corresponded with a rapid increase in temperatures and several freeze-thaw events (Figure 1), in comparison to a steady decline in germination for seeds put in the field in spring. In 1994–1995, germination of laboratory-stored seeds was up to 38% greater than seeds placed in the field, and germination of seeds placed in the field in spring was 7% to 61% greater than that of seeds put in the field in autumn. Germination of laboratory seeds was up to 95% greater than for seeds placed in the field in 1995–1996; seeds exposed to the spring environment germinated greater than ones also exposed to winter environments.

1998–1999

Water content of seeds, total germination, and seeds infected with fungi were affected ($P < 0.0001$) by the interacting influences of seed treatment and date of germination (Table 5). Water

TABLE 3

Water content of seeds and total germination for silver sagebrush during the seed retrieval test in 1995–1996

Seed treatment (location and time of placement)	Date of retrieval in 1995–1996							
	7 Nov	10 Apr	22 Apr	1 May	13 May	21 May	29 May	6 Jun
	Water content of seeds (% fresh weight)							
Laboratory	– ^a	17	16	9	6	5	11	4
Field—Autumn	–	31	29	24	8	48	13	7
Field—Spring	–	18	33	25	9	19	13	8
Standard error (date of germination X time of placement interaction)								4.5
	Total germination (%)							
Laboratory	79	82	71	81	83	74	82	84
Field—Autumn	78	65	52	56	58	49	52	46
Field—Spring	79	81	76	80	78	54	67	60
Standard error (date of germination X time of placement interaction)								4.8

^a Water content of seeds was not determined

TABLE 4

Levels of significance for linear, quadratic, and cubic contrasts of mean total germination over time for silver sagebrush seeds stored in the laboratory or placed in the field in autumn or spring on total germination

Year	Contrast	Linear	Quadratic	Cubic
1994–1995	Laboratory storage versus autumn and spring placement in the field	0.002	0.008	0.046
	Autumn versus spring placement in the field	0.00	0.019	0.025
1995–1996	Laboratory storage versus autumn and spring placement in the field	0.006	0.003	0.049
	Autumn versus spring placement in the field	0.805	0.272	0.041

content of seeds in field conditions varied considerably among dates of retrieval and averaged 34% compared to 15% for laboratory-stored seeds. Total germination of seeds stored in the laboratory varied between 82% and 94% while that of seeds placed in the field declined 81%. Germination for seeds placed in the field dropped sharply and the percentage of seeds infected with fungi increased dramatically as temperatures rose in March and many freezing and thawing events occurred (Figure 1). Of the seeds not germinating, 6% to 18% of laboratory-stored seeds were infected with fungi, but fungal infection increased from 12% to 83% in seeds exposed to field environments. The percentage of seeds (0.4%, $s_x = 0.12$) that stained with tetrazolium chloride was similar among dates ($P = 0.409$), seed treatments ($P = 0.682$) and the date X treatment interaction ($P = 0.369$).

Influence of Temperature on Germination in the Laboratory

Silver sagebrush seeds germinated over a broad range of temperatures (Figure 2). Germination was least at the extremes of the temperatures, and greatest at intermediate temperatures. Averaged over all 55-temperature combinations, total germination was 57% ($s_x = 2.4$) in 1994, 45% ($s_x = 2.0$) in 1995, and 53% ($s_x = 1.9$) in 1998.

DISCUSSION

Under natural situations, seeds of silver sagebrush are exposed to field environments of autumn through winter before germinating in spring and early summer (Romo and Grilz 2001). Silver sagebrush flowers from late August through early September (Budd and Campbell 1959), and most seeds mature and are dispersed in October (Romo personal observations) to early November (Wambolt and others 1989). Some seeds are maintained on mother plants into the following spring (Walton 1984; Romo personal observations). Variation in the rate of seed dispersal among individual plants of silver sagebrush (Wamboldt and others 1989) likely creates a gradient in time in

which seeds are exposed to environmental conditions in the seedbed. Seeds maintained in the canopy are not exposed as long to environmental conditions in the seedbed compared to seeds that are shed quickly from mother plants. A prolonged period of seed dispersal may enable some seeds of silver sagebrush to occupy safe sites in time similar to fringed sagebrush (Bai and Romo 1994, 1997) and Wyoming big sagebrush (Schuman and others 1998). Prolonged dispersal likely also enables some seeds to avoid stressful conditions of the seedbed.

Silver sagebrush seeds are not dormant (Walton 1984) and our findings indicate autumn-dispersed seeds could potentially germinate with cooling of

Figure 2 • Response surfaces for total germination of silver sagebrush seeds collected in 1994, 1995, and 1998 and incubated at constant and alternating temperatures ranging from 0 to 40 °C (32 to 104 °F).

TABLE 5

Water content of seeds, total germination, and non-germinating seeds infected with fungi when silver sagebrush seeds were exposed to laboratory or field conditions in 1998–1999

Seed treatment	Date of retrieval in 1998—1999								
(location)	17 Nov	23 Jan	20 Feb	20 Mar	31 Mar	24 Apr	14 May	28 May	21 Jun
	Water content of seeds (% fresh weight)								
Laboratory	20	18	18	15	16	13	12	14	13
Field	18	33	53	72	32	22	29	23	26
Standard error (location X date of germination)									2.7
	Total germination (%)								
Laboratory	89	84	92	93	88	82	91	87	94
Field	89	79	78	32	35	44	40	18	17
Standard error (location X date of germination)									5.5
	Non-germinating seeds infected with fungi (%)								
Laboratory	11	16	9	8	13	18	9	12	6
Field	12	20	22	68	65	56	60	82	83
Standard error (location X date of germination)									5.6

the environment. Late-autumn dispersal coupled with temperatures that are unlikely to be warm enough or of adequate duration to permit germination until the following spring reduces the probability of germination in autumn or winter to near nil. It is possible, however, that seeds could germinate in autumn if temperatures and soil water are adequate when artificially seeded in early autumn. Seedlings emerging in autumn would be predisposed to freezing temperatures of autumn and winter, with 100% mortality expected for seedlings at temperatures below -13°C (9°F) (Hou and Romo 1998b). Winter temperatures around -40°C (-40°F) are common in our study area (Environment Canada 1982).

If conditions are not conducive to germination in autumn, silver sagebrush seeds would be exposed to field environments over the winter before germinating the following spring or summer. Reduced germination and viability, with greater reductions occurring during autumn- than in spring-placed seeds indicates the field environment, particularly during the winter, is stressful to many seeds in the population studied. This reduction in germination contrasts with improved germination after exposure to field environments in shrubs with dormant seeds (Haferkamp and others 1990; Shaw and others 1994). Bewley and Black (1982) argued that seeds could be damaged by freezing temperatures if their water content is greater than about 14%. Freezing damage to seeds is probable in our studies because water content exceeded 14% in both autumn- and spring-placed seeds, and seeds were exposed to several freezing

events, particularly in spring, with more occurring in the former treatment.

The decline of germination as temperatures rose in spring is consistent with the prediction that aging of moist seeds increases with temperature (Ellis and Roberts 1980). Increasing temperatures, precipitation, and relative humidity reduces germination of soybean (*Glycine max* (L.) Merr. [Fabaceae]) (TeKrony and others 1980). Exposing seeds to unfavorable environmental conditions for longer periods reduces germination more than exposure for shorter periods (Roberts and Ellis 1982). Greater reduction in germination for autumn-placed seeds is attributed to longer exposure to more unfavorable environmental conditions than seeds put out in spring.

Nearly all of the seeds that failed to germinate were not viable. More than 90% of big sagebrush (*Artemisia tridentata* Nutt.) seeds lost viability within 6 mo, and 100% decomposed within 10 mo of being placed in soil (Crist and Friese 1993). Young and Evans (1989) concluded that big sagebrush (*Artemisia tridentata* Nutt. ssp. *tridentata*) does not develop a soil seed bank. Anecdotal evidence prompted Schuman and others (1998) to conclude that seeds of Wyoming big sagebrush may enter dormancy and persist in the soil. Fringed sagebrush, a common associate with silver sagebrush in the Northern Mixed Prairie (Coupland 1950), maintains a persistent soil seed-bank (Bai and Romo 1994, 1996). Even though germination of silver sagebrush declined with exposure to field environments, some seeds remained viable, suggesting that a soil seedbank

could develop for this shrub. Apparently no studies have examined the dynamics of silver sagebrush seed reserves in soil.

Seeds dispersed in autumn through spring can germinate as temperatures rise in spring and early summer. As predicted from our germination tests, most seedlings of silver sagebrush emerge in late spring and early summer (Romo and Grilz 2001). Romo and Eddleman (1995) also concluded that seeds and seedlings of silver sagebrush are adapted to germinate and grow at low temperatures. Increasing germination with warming temperatures presumably improves the probability that most seeds of silver sagebrush germinate early in the growing season, maximizing the period for seedling growth in this environment with a short growing season. Early emerging seedlings are also more vigorous than later emerging ones (Hou and Romo 1998a). Conversely, freezing temperatures or desiccation may kill seedlings emerging early in spring (Hou and Romo 1997, 1998b).

The fact that some seeds did not germinate at the lowest temperatures tested in the laboratory suggests that additional seeds could germinate as temperatures warm later in spring and summer. Stratification has no effect on germination of silver sagebrush (Walton 1984). Later-emerging seedlings might avoid freezing temperatures of early spring, however, they would face an abbreviated period for growth, especially because of soil water limitations during summer. Summer drought is a major cause of seedling death in silver sagebrush and more seedlings establish in wet than dry years (Walton 1984). Later emerging seedlings are less tolerant of freezing temperatures than early emerging ones (Hou and Romo 1998b). Therefore, although later emerging seedlings can avoid freezing temperatures in spring, they may be predisposed to summer drought or freezing damage the following winter.

SUMMARY

Our studies were conducted on seeds from a population at the northern reaches of the distribution of silver sagebrush. Although ecotypic studies have yet to be completed for silver sagebrush, seeds of ecotypes from different environments may have adaptive germination responses that are unique as shown for big sagebrush, Wyoming big sagebrush, and mountain big sagebrush (McDonough and Harniss 1974; Meyer and others 1990; Meyer and Monsen 1991, 1992). Silver sagebrush seeds germinate over a broad range of temperatures like other species of sagebrush (Young and others 1991; Bai and others 1995). Viability in a substantial portion of the seed population declines with increasing exposure to environmental conditions in the field. Seeds put in the field in autumn experience stress-

ful environmental conditions longer and hence germination is reduced more than spring-placed seeds. High water content and increasing temperatures over time (Ellis and Roberts 1980; Roberts and Ellis 1982) likely caused the decline in germination under field environments. Despite seed viability declining under field environments, 19% to 82% of the seeds remained viable, suggesting that silver sagebrush may develop a soil seedbank. Greater contributions to the seedbank are predicted for seeds that are naturally dispersed or artificially sown in spring versus autumn. Our studies strongly implicate loss of seed viability in the field as a primary factor limiting establishment (Eddleman 1980; Walton 1984; Romo and Grilz 2001) of this shrub from seeds. Since most seedlings of silver sagebrush emerge in spring and early summer (Romo and Grilz 2001), seeding in early spring is recommended to maintain high viability and germination of seeds in this shrub.

REFERENCES

- Allen MF. 1988. Belowground structure: a key to reconstructing a productive arid ecosystem. In: Allen EB, editor. The reconstruction of disturbed arid lands—an ecological approach. Boulder (CO): Westview Press. p 113–135.
- Bai Y, Romo JT. 1994. Germination of previously buried seeds of fringed sagebrush (*Artemisia frigida*). *Weed Science* 42:390–397.
- Bai Y, Romo JT. 1996. Fringed sagebrush response to sward disturbances: seedling dynamics and plant growth. *Journal of Range Management* 49:228–233.
- Bai Y, Romo JT. 1997. Seed production, seed rain, and the seedbank of fringed sagebrush. *Journal of Range Management* 50:151–155.
- Bai Y, Romo JT, Young JA. 1995. Influences of temperature, light and water stress on germination of fringed sagebrush (*Artemisia frigida*). *Weed Science* 43:219–225.
- Bai Y, Booth DT, Roos EE. 1997. Effect of seed moisture on Wyoming big sagebrush seed quality. *Journal of Range Management* 50:419–422.
- Barker WT, Whitman WC. 1988. Vegetation of the northern Great Plains. *Rangelands* 10:266–272.
- Beetle AA. 1960. A study of sagebrush: the section *Tridentatae* of *Artemisia*. Laramie (WY): University of Wyoming Agriculture Experiment Station. Bulletin No. 368. 83 p.
- Bewley JD, Black M. 1982. Physiology and biochemistry of seeds in relation to germination. Volume 2. New York (NY): Springer-Verlag. 375 p.
- Budd AC, Campbell JB. 1959. Flowering sequence of a local flora. *Journal of Range Management* 12:127–132.
- Call CA, Roundy BA. 1991. Perspectives and processes in revegetation of arid and semi-arid rangelands. *Journal of Range Management* 44:543–549.
- Coupland RT. 1950. Ecology of mixed prairie in Canada. *Ecological Monographs* 20:273–315.
- Coupland RT. 1961. A reconsideration of grassland classification in the Northern Great Plains of North America. *Journal of Ecology* 49:135–167.
- Coupland RT. 1992. Mixed prairie. In: Coupland RT, editor. *Natural grasslands-Introduction and Western Hemisphere. Ecosystems of the World 8A*. New York (NY): Elsevier Science Publishers. p 151–182.
- Crist TO, Friese CF. 1993. The impact of fungi on soil seeds: implications for plants and granivores in a semi-arid shrub steppe. *Ecology* 74:2231–2239.

- Eddleman, LE. 1980. Coal mine reclamation with native plants. In: Symposium proceedings, symposium on watershed management; 1980 Jul 21–30; Boise, ID. New York (NY): American Society of Civil Engineers. p 80–90.
- Ellis RH, Roberts EH. 1980. The influence of temperature and moisture on seed viability in barley (*Hordeum distichum* L.) Annals of Botany 45:31–37.
- Environment Canada, Atmospheric Environmental Service. 1982. Canadian climate normals (1951–1980), temperature and precipitation (Prairie provinces). Ottawa (ON). 266 p.
- Evans RA, Easi DA, Book DN, Young JA. 1982. Quadratic response surface analysis of seed germination trials. Weed Science 30:411–416.
- Grabe DL. 1972. Tetrazolium testing handbook for agricultural seeds. Contribution 29, handbook on seed testing. Springfield (IL): Association of Official Seed Analysts. 62 p.
- Haferkamp MR, Ganskopp DC, Marietta KL, Knapp BW. 1990. Environmental influences on germination of utricles and seedling establishment of immigrant forage kochia. Journal of Range Management 43:518–522.
- Hanson HC, Whitman W. 1938. Characteristics of major grassland types in western North Dakota. Ecological Monographs 8:59–114.
- Harniss RO, McDonough WT. 1976. Yearly variation in germination in three subspecies of big sagebrush. Journal of Range Management 29:167–168.
- Harvey SJ. 1981. Life history and reproductive strategies in *Artemisia* [MSc thesis]. Bozeman (MT): Montana State University. 132 p.
- Hegarty TW. 1978. The physiology of seed hydration and dehydration, and the relation between water stress and the control of germination: a review. Plant, Cell, and Environment 1:101–119.
- Hou JQ, Romo JT. 1997. Growth and freezing tolerance of winterfat seedlings. Journal of Range Management 50:165–169.
- Hou JQ, Romo JT. 1998a. Seed weight and germination time affect growth of two shrubs. Journal of Range Management 51:699–703.
- Hou JQ, Romo JT. 1998b. Cold-hardiness of silver sagebrush seedlings. Journal of Range Management 51:704–708.
- Hulett GK, Coupland RT, Dix RL. 1966. The vegetation of dune and sand areas within the grassland region of Saskatchewan. Canadian Journal of Botany 44:1307–1331.
- Lauenroth WK, Milchunas DG. 1992. Shortgrass steppe. In: Coupland RT, editor. Natural grasslands—Introduction and Western Hemisphere. Ecosystems of the World 8A. New York (NY): Elsevier Science Publishers. p 183–226.
- Lawrence DL, Romo JT. 1994. Tree and shrub communities of wooded draws near the Matador Research Station in southern Saskatchewan. Canadian Field-Naturalist 108:397–412.
- Looman J, Best KF. 1979. Budd's flora of the Canadian Prairie Provinces. Hull (QC): Canadian Government Publishing Centre. Supply and Services Canada. 863 p.
- McDonough WT, Harniss RO. 1974. Effects of temperature on germination in three subspecies of big sagebrush. Journal of Range Management 27:204–205.
- Meyer SE, Monsen SB. 1991. Habitat-correlated variation in mountain big sagebrush (*Artemisia tridentata* ssp. *vaseyana*) seed germination patterns. Ecology 72:739–742.
- Meyer SE, Monsen SB. 1992. Big sagebrush germination patterns: subspecies and population differences. Journal of Range Management 4:87–93.
- Meyer SE, Monsen SB, McArthur ED. 1990. Germination response of *Artemisia tridentata* (Asteraceae) to light and chill: patterns of between-population variation. Botanical Gazette 151:176–183.
- Miller RM. 1987. Mycorrhizae and succession. In: Jordan III WR, Gilpin ME, Aber JD, editors. Restoration ecology—a synthetic approach to ecological research. New York (NY): Cambridge University Press. p 205–219.
- Petersen RG. 1985. Design and analysis of experiments. New York (NY): Marcel Dekker Inc. 429 p.
- Pyke DA, Archer S. 1991. Plant-plant interactions affecting plant establishment and persistence on revegetated rangeland. Journal of Range Management 44: 550–557.
- Roberts EH, Ellis RH. 1982. Physiological, ultrastructural and metabolic aspects of seed viability. In: Khan AA, editor. The physiology and biochemistry of seed development, dormancy and germination. New York (NY): Elsevier Biomedical Press. p. 465–485.
- Romo JT, Eddleman LE. 1995. Use of degree-days in multiple-temperature experiments. Journal of Range Management 48:410–416.
- Romo JT, Grilz RW. 2002. Establishment of silver sagebrush in the Northern Mixed Prairie. Journal of Range Management 55. Forthcoming.
- Roos EE. 1986. Precepts of successful seed storage. In: McDonald MB, Nelson C, editors. Physiology of seed deterioration. Madison (WI): Crop Science Society of America Special Publication No. 11. p 1–25.
- Rowe JS, Coupland RT. 1984. Vegetation of the Canadian prairies. Prairie Forum 9:231–248.
- Schuman GE, Booth DT, Cockrell, JR. 1998. Cultural methods for establishing Wyoming big sagebrush on mined lands. Journal of Range Management 51:223–230.
- Shaw NL, Haferkamp MR, Hurd EG. 1994. Germination and establishment of spiny hopsage in response to planting date and seedbed environment. Journal of Range Management 47:165–174.
- Snedecor GW, Cochran WG. 1980. Statistical methods. 7th ed. Ames (IA): Iowa State University Press. 507 p.
- TeKrony DM, Egli DB, Phillips AD. 1980. Effect of field weathering on the viability and vigor of soybean seed. Agronomy Journal 72:749–753.
- Walton TP. 1984. Reproductive mechanisms of plains silver sagebrush *Artemisia cana cana* in southern Montana. [MSc thesis]. Bozeman (MT): Montana State University. 161 p.
- Wambolt CL, Walton TP, White RS. 1989. Seed dispersal characteristics of plains silver sagebrush. Prairie Naturalist 21:113–118.
- Welch BL. 1996. Effects of humidity on storing big sagebrush seed. Ogden (UT): USDA Forest Service, Intermountain Research Station. Research Paper INT–RP–493. 5 p.
- Welch BL, Briggs SF, Johansen JH. 1996. Big sagebrush seed storage. Ogden (UT): USDA Forest Service, Intermountain Research Station. Research Note INT–RN–430. 3 p.
- Young JA, Evans RA. 1989. Dispersal and germination of big sagebrush (*Artemisia tridentata*) seeds. Weed Science 7:201–206.
- Young JA, Palmquist DE, Evans RA. 1991. Temperature profiles for germination of big sagebrush seeds from native stands. Journal of Range Management 44:385–390.

AUTHOR INFORMATION

James T Romo
Professor
Department of Plant Sciences
51 Campus Drive
University of Saskatchewan
Saskatoon, SK S7N 5A8
romo@sask.usask.ca

James A Young
Range Scientist
USDA Agriculture Research Service
920 Valley Road
Reno, NV 89512
jayoung@scs.unr.edu