



REFEREED RESEARCH

- Evaluating the use of woody debris to enhance native plant establishment from seeds on legacy coal mines in West Virginia (USA) 272
Anna M Branduzzi, Christopher D Barton, Carol C Baskin, and Allison G Davis
- Cutting propagation of 4 Florida native taxa of wild coffee (*Psychotria* spp.) 288
Teagan H Young, Sandra B Wilson, Mack Thetford, and James Colee
- Seedling emergence and seed production of curlycup gumweed 299
Derek Tilley, Mary Wolf, Derek Jolley, and Gordon Hirning

PROPAGATION PROTOCOL

- Germination and propagation of *Lithospermum canescens* (Michx.) Lehm., a difficult-to-grow prairie species 309
David Sollenberger, Cathy Thomas, and Louise Egerton-Warburton

GENERAL TECHNICAL

- Hypersaturated Piedmont prairies: a screening process for optimum placement 324
Sean Dunlap, Brad Davis, and Jon Calabria
- Propagation for commercial production of sweet acacia (*Vachellia farnesiana*): a native plant with ornamental potential 337
Thomas P Smith IV, Sandra B Wilson, S Christopher Marble, and Jianjian Xu
- Restoration of a calcareous river floodplain in southwestern Montana, part 2: presentation of monitoring data 349
Tara Luna

AUTHOR GUIDELINES 362

Photo credits: (top) Polycotyledonous behavior of sweet acacia (*Vachellia farnesiana* (L.) Wight & Arn. [Fabaceae]) seedlings, photo by Thomas Smith IV; (middle) seeding depth evaluation for curlycup gumweed (*Grindelia squarrosa* (Pursh) Dunal [Asteraceae]) in 3 soil textures (fine sand, loam, and clay loam), photo by Derek Tilley; (bottom) research plots in a red spruce ecosystem (*Picea rubens* Sarg. [Pinaceae]) of West Virginia examine direct seeding of native species, with and without woody debris amendments, photo by Christopher Barton.